

Standard Anterior Cruciate Ligament Reconstruction Rehabilitation Protocol

Time	Goals	Precautions/Restrictions	Treatment
Weeks 0 – 2	<ul style="list-style-type: none"> • Protect surgical site • Reduce pain and swelling • Active ROM: 0-90 degree • Full passive extension • Active quadriceps control • Reduce muscle atrophy • Safe use of crutches with near normal gait mechanics 	<ul style="list-style-type: none"> • No brace use • WBAT with crutches • No resisted open chain knee extensions for six weeks • No isolated hamstring strengthening for hamstring autograft surgery 	<ul style="list-style-type: none"> • ROM (as tolerated) <ul style="list-style-type: none"> ○ PROM – AAROM - AROM ○ Patella mobilizations • Quadriceps recruitment/NMES • Global LE/hip strengthening • Gait training with crutches • Modalities as indicated <ul style="list-style-type: none"> ○ Cryotherapy: 5-7 times per day • Initial Visit: FOTO, LEFS, PSFS • 2 week follow up with MD
Weeks 2 – 6	<ul style="list-style-type: none"> • Full, symmetric and pain-free AROM • SLR without extensor lag • Normalized gait mechanics • DL squat with good mechanics • Progression of quadriceps strength/endurance • Increase functional activities 	<ul style="list-style-type: none"> • Open kinetic chain is limited to bodyweight leg extensions (weeks 2-6) • No resisted open kinetic chain exercises • Initiate isolated hamstring strengthening for hamstring autograft surgery (6 weeks) • No running, jumping, cutting, pivoting, or twisting • Avoid painful activities/exercises 	<ul style="list-style-type: none"> • AAROM - AROM • Gait training progressing from assistive device • Core stabilization exercises • Closed kinetic chain strengthening as tolerated • Global LE strengthening <ul style="list-style-type: none"> ○ Begin functional strengthening exercises (bridge, mini-squat, step up, etc) • Double limb to single limb balance/proprioception • Aerobic training: <ul style="list-style-type: none"> ○ Walking program when walking with normal gait mechanics ○ Stationary bike • Optional therapies (if available/as indicated): <ul style="list-style-type: none"> ○ BFR therapy ○ Anti-gravity treadmill for walking gait ○ Aquatic therapy once incision is healed and cleared by surgeon (2-4 weeks) ○ NMES • Modalities as indicated • Week 6: FOTO, LEFS
Weeks 6 – 12	<ul style="list-style-type: none"> • Full, symmetric and pain-free ROM • Progress quadriceps strength/endurance • Increase functional activities 	<ul style="list-style-type: none"> • May initiate resisted open kinetic chain exercise <ul style="list-style-type: none"> ○ 90-45° at 6 weeks ○ 90-30° at 8 weeks ○ 90-0° at 10 weeks ○ 90-0° with progressive loading at 12 weeks • No running, jumping, cutting, pivoting, or twisting • Avoid painful activities/exercises • Avoid patellofemoral pain 	<ul style="list-style-type: none"> • End range flexion and extension • Aerobic training on stationary bike, elliptical, stair climber, UBE • Core stabilization exercises • Progressive double and single limb strengthening • Single to multi-plane exercise • Progression of balance/proprioception • Modalities as indicated • Week 12: FOTO, LEFS, PSFS, ACL-RSI • 6-8 week follow up with MD (SGYM)

This protocol is not meant to be prescriptive but a recommendation to guide the rehabilitation process.

Each patient’s progress may vary based on specifics of their injury and procedure.

<p>Weeks 12-16</p>	<ul style="list-style-type: none"> • Full, symmetric ROM • No effusion with increased activity • Increase intensity and duration of functional LE strength • Initiate return to jogging program • Begin low level plyometric and agility training 	<ul style="list-style-type: none"> • Avoid painful activities/exercises • Jogging program initiated at 12 weeks if cleared by surgeon <ul style="list-style-type: none"> ○ No effusion ○ Full AROM ○ >80% LSI • No jogging on painful or swollen knee 	<ul style="list-style-type: none"> • Increase loading capacity for lower extremity strengthening exercises • Continue balance/proprioceptive training • Week 12: begin return to jogging program <ul style="list-style-type: none"> ○ If applicable, start with pool/anti-gravity treadmill • Begin low level plyometric and agility training at 12 weeks • Functional assessment (see attached) • 3-4 month follow up with MD (SGYM)
<p>Months 4-6</p>	<ul style="list-style-type: none"> • Continue to progress functional strengthening • Successful progression of the return to running program • Initiate higher level plyometric and agility training 	<ul style="list-style-type: none"> • No jogging on a painful or swollen knee • Avoid painful activities/exercises • Avoid patellofemoral pain • No participation in sports unless specified by care team 	<ul style="list-style-type: none"> • Progression of return to jogging program • Gradually increase lifting loads focusing on form, control, and tissue tolerance • Progress as tolerated: <ul style="list-style-type: none"> ○ Core Stability ○ Strength ○ Endurance ○ Proprioception/Balance • Increase intensity of plyometric and agility training • Foot speed and change of direction • Functional assessment at 6 months (see attached) • Month 6: FOTO, LEFS, PSFS, ACL-RSI
<p>Months 6 – 9</p>	<ul style="list-style-type: none"> • Continue to progress functional strengthening • Sport-specific training 	<ul style="list-style-type: none"> • No participation in sports unless specified by your care team • Avoid painful activities 	<ul style="list-style-type: none"> • Progress as tolerated: <ul style="list-style-type: none"> ○ Core Stability ○ Strength ○ Endurance ○ Proprioception/Balance • Begin sport-specific training • Single-to multi-task • Reactionary drills • Perturbation training • Closed to open environment
<p>Months 9+</p>	<ul style="list-style-type: none"> • Pass return to play criteria (re-test at 12+ months, if necessary) • Begin gradual return to sport 	<ul style="list-style-type: none"> • Gradual return to full participation in sports 	<ul style="list-style-type: none"> • Progress as tolerated: <ul style="list-style-type: none"> ○ Core Stability ○ Strength ○ Endurance ○ Proprioception/Balance ○ Plyometric training ○ Agility drills ○ Sport-specific activities • Gradual return to sport progression • Month 9: FOTO, LEFS, PSFS, ACL-RSI • Functional assessment (see attached) • 9+ month follow up with MD (SGYM)

This protocol is not meant to be prescriptive but a recommendation to guide the rehabilitation process. Each patient's progress may vary based on specifics of their injury and procedure.

Anterior Cruciate Ligament Testing Protocol

Phase	Goals	Surgery	Testing
Week 12 (SGYM)	<ul style="list-style-type: none"> • Full, symmetric ROM • Y-balance anterior reach asymmetry < 5 cm • Quadriceps strength for isometric test > 80% of uninvolved side 	<ul style="list-style-type: none"> • ACL reconstruction • ACL reconstruction with meniscus repair • ACL Allograft (12 week and 6 month recheck) • ACL revision (12 week and 6 month recheck) 	<p>TESTING:</p> <ul style="list-style-type: none"> • Knee assessment including assessment for effusion • Passive and active ROM • Y-balance anterior reach • Isometric knee extension at 60° and 90° • FOTO, LEFS, PSFS, ACL-RSI
Month 6 (No SGYM)	<ul style="list-style-type: none"> • Full, symmetric ROM • Y-balance anterior reach asymmetry < 3 cm • > 80% LSI for isokinetic testing • 90% LSI for functional testing • ACL-RSI > 56 	<ul style="list-style-type: none"> • ACL reconstruction • ACL reconstruction with meniscus repair • <u>ACL Allograft (9 month)</u> • <u>ACL revision (9 month)</u> 	<p>TESTING:</p> <ul style="list-style-type: none"> • Knee assessment including assessment for effusion • Passive and active ROM • Hop Test <ul style="list-style-type: none"> ○ Single Hop ○ Triple Hop • Isokinetic Test (90°, 180°/s) • Y-Balance Anterior Reach • Agility T-Test • FOTO, LEFS, PSFS, ACL-RSI
Month 9-10 (SGYM)	<ul style="list-style-type: none"> • Full, symmetric ROM • > 90% LSI for isokinetic and functional testing • Y-balance anterior reach asymmetry < 3 cm • ACL-RSI > 56 	<ul style="list-style-type: none"> • ACL reconstruction • ACL reconstruction with meniscus repair • <u>ACL Allograft (12 month)</u> • <u>ACL revision (12 month)</u> 	<p>TESTING:</p> <ul style="list-style-type: none"> • Knee assessment including assessment for effusion • Passive and active ROM • Hop Test <ul style="list-style-type: none"> ○ Single Hop ○ Triple Hop ○ Cross-over Hop • Isokinetic Test (90°, 180°/s) • Y-Balance Anterior Reach • Agility T-Test • FOTO, LEFS, PSFS, ACL-RSI

This protocol is not meant to be prescriptive but a recommendation to guide the rehabilitation process.
Each patient's progress may vary based on specifics of their injury and procedure.